

## EXTRAGONADAL GERM CELL TUMOR

### Procedure

- Marginal resection
- Wide resection
- Other: (specify)

### Patient Age

- Congenital/neonatal (birth – 6 mo)
- Childhood/prepubertal (7 mo – 11 y)
- Postpubertal/adult ( $\geq 12$  y)

### Tumor Site

- Intracranial
- Head and neck region (excluding intracranial)
- Mediastinum
- Retroperitoneum/Abdomen
- Sacrococcygeal
- Other: (specify)

### Tumor Size

Greatest dimension: \_\_\_ cm

### Histologic type (select all that apply)

- Mature teratoma
- Immature teratoma
- Mature or immature teratoma with additional malignant somatic component: (specify type, eg, epithelial malignancy, sarcoma)
- Germinoma
- Yolk sac tumor
- Embryonal carcinoma
- Choriocarcinoma
- Mixed germ cell tumor

### Histologic Grade (applicable to immature teratomas only)

- Grade 1
- Grade 2
- Grade 3
- Percent of teratoma composed of immature elements: \_\_\_% (if applicable)

**Treatment History**

- No known preoperative therapy
- Preoperative therapy given
- Not specified

**Treatment Effect**

- Not applicable (no known preoperative therapy)
- Not identified
- Present; percent tumor necrosis: \_\_\_%

**Microscopic Tumor Extension (applicable to sacrococcygeal resections only)**

- Tumor involving coccyx identified
- Coccyx uninvolved

**Margins**

- Cannot be assessed
- Uninvolved by tumor (closest margin and distance from closest margin: \_\_\_ , \_\_\_ cm)
- Involved by tumor; margin(s) involved: \_\_\_

**Lymphovascular Invasion**

- Not identified
- Present

**Perineural Invasion**

- Not identified
- Present

**Regional Lymph Nodes**

- No nodes submitted or found
- Number of lymph nodes involved/total number of lymph nodes submitted: \_\_\_/\_\_\_
- Sites of involved lymph nodes (if known) and histologic type(s) present:

**Distant Metastasis**

- Not applicable (no distant sites submitted)
- Present