

GALLBLADDER CARCINOMAS

Procedure

Cholecystectomy
Radical cholecystectomy (with liver resection and lymphadenectomy)
Other (specify): _____

Tumor Site (select all that apply)

Fundus.
Body.
Neck.
Cystic duct.
Free peritoneal side of gallbladder.
Hepatic side of gallbladder.
Cannot be determined.

Tumor Size: ____ cm

Histologic Type

Adenocarcinoma
Mucinous adenocarcinoma
Other (specify): _____

Histologic Grade

Well differentiated, G1
Moderately differentiated, G2
Poorly differentiated, G3
Undifferentiated

Tumor Extension

Invades lamina propria
Invades muscular layer
Invades perimuscular connective tissue on peritoneal side without serosal involvement
Invades perimuscular connective tissue on hepatic side without liver involvement
Perforates serosa (visceral peritoneum)
Invades liver
Invades adjacent organs and/or structures (specify)#: _____
Invades main portal vein and/or hepatic artery
No evidence of primary tumor

Adjacent organs/structures include stomach, duodenum, colon, pancreas, extrahepatic bile ducts, omentum, main portal vein, and hepatic artery

Margins

- Negative (all margins including cystic duct margin are uninvolved by invasive carcinoma or high-grade biliary intraepithelial neoplasia)
- Negative (all margins including cystic duct and hepatic parenchymal margins are uninvolved by invasive carcinoma or high-grade biliary intraepithelial neoplasia)
- All margins negative for invasive carcinoma. Cystic duct margin involved by high-grade biliary intraepithelial neoplasia.
- Positive (specify margin):

Regional Lymph Nodes

No lymph nodes submitted or found

Number of positive lymph nodes / Total number of nodes examined: ____/____

Distant Metastasis: Yes (specify sites): Unknown

Pathologic Stage Classification (AJCC 8th Edition)

Note: Modify the pTNM stage with **m** (multiple primaries - suffix), **r** (recurrent) and/or **y** (post-treatment) as applicable

T stage

T0: No evidence of primary tumor

T1a: Tumor invades lamina propria

T1b: Tumor invades muscular layer

T2a: Tumor invades perimuscular connective tissue on peritoneal side, without involvement of serosa

T2b: Tumor invades the perimuscular connective tissue on hepatic side, with no extension into liver

T3: Tumor perforates serosa and/or directly invades liver and/or **one** other adjacent organ or structure

T4: Tumor invades main portal vein and/or hepatic artery OR invades **two or more** extrahepatic organs or structures

N stage

N0: No regional lymph node metastasis

N1: Metastases to one to three regional lymph nodes

N2: Metastases to four or more regional lymph nodes

M stage

M1: Distant metastasis