

LARYNX NEOPLASM (GLOTTIS)

Procedure: Total laryngectomy
Other: _____

Tumor site: Glottis
Other: _____

Tumor Laterality: Right
Left
Midline
Bilateral
Other: _____

Tumor Focality: Unifocal Multifocal

Tumor Size (greatest dimension): _____ cm

Histologic Type: Keratinizing squamous cell carcinoma
Nonkeratinizing squamous cell carcinoma
Papillary squamous cell carcinoma
Adenosquamous carcinoma
Basaloid squamous cell carcinoma
Spindle cell (sarcomatoid) squamous cell carcinoma
Verrucous carcinoma
Other: _____

Histologic Grade: Well-differentiated (G1)
Moderately-differentiated (G2)
Poorly-differentiated (G3)
Other: _____

Lymphovascular Invasion: Present Absent

Perineural Invasion: Present Absent

Cartilage Invasion: Present Absent N/A

Bone Invasion: Present Absent N/A

Margins: Negative Positive

Extent of Primary Tumor:

- No evidence of primary tumor (pT0)
- Carcinoma in situ (pTis)
- Tumor limited to one vocal cord(s) (may involve anterior or posterior commissure) with normal mobility (T1a)

- Tumor involves both vocal cords (may involve anterior or posterior commissure) with normal mobility (T1b)
- Tumor extends to supraglottis and/or subglottis and/or with impaired vocal cord mobility (pT2)
- Tumor limited to the larynx with vocal cord fixation and/or invades paraglottic space and/or inner cortex of the thyroid cartilage (pT3)
- Tumor invades through the outer cortex of the thyroid cartilage and/or invades tissues beyond the larynx (eg, trachea, soft tissues of neck including deep extrinsic muscle of the tongue, strap muscles, thyroid, or esophagus) (pT4a)
- Tumor invades prevertebral space, encases carotid artery, or invades mediastinal structures (pT4b)

Lymph Node Status:

- Not assessed
- No regional lymph node metastasis (pN0)
- Metastasis in a single ipsilateral lymph node, 3 cm or smaller in greatest dimension without extranodal extension (pN1)
- Metastasis in a single ipsilateral lymph node, 3 cm or smaller in greatest dimension with extranodal extension (pN2a)
- Metastasis in a single ipsilateral lymph node, larger than 3 cm but not larger than 6 cm in greatest dimension without extranodal extension (pN2a)
- Metastasis in multiple ipsilateral lymph nodes, none larger than 6 cm in greatest dimension without extranodal extension (pN2b)
- Metastasis in bilateral or contralateral lymph nodes, none larger than 6 cm in greatest dimension without extranodal extension (pN2c)
- Metastasis in a lymph node larger than 6 cm in greatest dimension without extranodal extension (pN3a)
- Metastasis in a single ipsilateral lymph node, larger than 3 cm in greatest dimension with extranodal extension (pN3b)
- Metastases in multiple ipsilateral, contralateral, or bilateral lymph nodes with extranodal extension (pN3b)
- Metastasis in a single contralateral node, 3 cm or smaller with extranodal extension (pN3b)

Extent of Primary Tumor (Mucosal Melanoma):

- *Tumors limited to the mucosa and immediately underlying soft tissue, regardless of thickness or greatest dimension (pT3)*
- *Tumor involving deep soft tissue, cartilage, bone, or overlying skin (pT4a)*
- *Tumor involving brain, dura, skull base, lower cranial nerves (IX, X, XI, XII), masticator space, carotid artery, prevertebral space, or mediastinal structures (pT4b)*

Lymph Node Status (Mucosal Melanoma):

- *Not assessed*
- *No regional lymph node metastasis (pN0)*
- *Regional lymph node metastases present (pN1)*

Distant Metastasis:

- Not assessed
- Distant metastasis (pM1)