

Pleural Mesothelioma: Resections ONLY (not intended for biopsies)

Specimen laterality:

- RIGHT or LEFT Pleura
- Other (specify)

Procedure:

- Extended pleurectomy/decortication
- Pleurectomy/decortication
- Partial pleurectomy
- Extrapleural pneumonectomy
- Other (specify)
- Not specified

Sites involved by tumor/Tumor extent (select all that apply):

- Parietal pleura
- Visceral pleura
- Diaphragm
- Other (specify)
- Not specified

Tumor focality:

- Diffuse
- Localized (sharply circumscribed)
- Cannot be determined

†Tumor size (localized tumors only)

___ cm x ___ cm x ___ cm

Histologic type:

- Epithelioid mesothelioma
- Sarcomatoid mesothelioma
- Desmoplastic mesothelioma
- Biphasic mesothelioma
- Other (specify)

Tumor extension (select all that apply)

- Limited to parietal pleura (NEGATIVE for ipsilateral visceral, mediastinal, or diaphragmatic involvement)
- Limited to parietal pleura WITH FOCAL involvement of ipsilateral visceral, mediastinal, or diaphragmatic pleura
- Tumor involves all ipsilateral pleural surfaces (including fissure)
- Tumor involves diaphragmatic muscle
- Tumor extends into lung parenchyma
- Tumor involves endothoracic fascia

Tumor extends into mediastinal fat
Solitary focus extends into the soft tissues of the chest wall
Diffuse or multiple foci extend into soft tissue of chest wall
Tumor extends into but not through the pericardium
Tumor involves rib(s)
Tumor involves mediastinal organ(s) (specify):
Other (specify):
Cannot be assessed

Margins:

Not applicable
Cannot be assessed
Negative
Involved by mesothelioma (specify)

Regional lymph nodes

No lymph nodes submitted or found

Number of lymph nodes involved ____
Number cannot be determined (explain):
Number of lymph nodes examined ____
Number cannot be determined (explain):

Treatment effect:

Not applicable
More than 50% residual viable tumor
Less than 50% residual viable tumor
Cannot be determined

Pathologic stage (TNM; AJCC 8th ed)