

RETROPERITONEAL LYMPH NODE DISSECTION FOR METASTATIC GERM CELL TUMOR

Histologic Viability of Tumor:

| | | | |
|----------------------------------|---------|--------|-----|
| Necrosis: | Present | Absent | N/A |
| Viable tumor: | Present | Absent | N/A |
| Viable teratoma: | Present | Absent | N/A |
| Viable non-teratomatous tumor: | Present | Absent | N/A |
| Types of non-teratomatous tumor: | | | |
| Seminoma: | Present | Absent | N/A |
| Embryonal carcinoma: | Present | Absent | N/A |
| Yolk sac tumor: | Present | Absent | N/A |
| Choriocarcinoma: | Present | Absent | N/A |
| Other: | | | |

Lymph node number (fraction involved by tumor): _____ / _____

Site(s) of lymph nodes involved by tumor: _____

Size of the largest lymph node involved by tumor: _____ cm

Extranodal extension: Present Absent N/A

Lymph node status:

- No regional lymph node metastasis (pN0)
- Metastasis with a lymph node mass 2 cm or smaller in greatest dimension and less than or equal to 5 nodes positive, none larger than 2 cm in greatest dimension (pN1)
- Metastasis with a lymph node mass larger than 2 cm but smaller than 5 cm in greatest dimension; or more than 5 nodes positive, none larger than 5 cm; or evidence of extranodal extension of tumor (pN2)
- Metastasis in a lymph node larger than 5 cm in greatest dimension (pN3)